International Journal of Research in Agronomy

E-ISSN: 2618-0618 P-ISSN: 2618-060X © Agronomy www.agronomyjournals.com 2023; 6(1): 01-05 Received: 07-10-2022 Accepted: 10-11-2022

Navjot Singh Assistant Professor, Department of Agriculture, GSSDGS Khalsa College, Patiala, Punjab, India

Mohinder Lal

Assistant Professor, Department of Agriculture, GSSDGS Khalsa College, Patiala, Punjab, India

Corresponding Author: Mohinder Lal Assistant Professor, Department of Agriculture, GSSDGS Khalsa College, Patiala, Punjab, India

Response of different weed management practices on growth and yield of black gram (*Vigna mungo* L.) under irrigated conditions of Punjab

Navjot Singh and Mohinder Lal

DOI: https://doi.org/10.33545/2618060X.2023.v6.i1a.154

Abstract

The field experiment was carried out during the summer season of 2021-2022 to study the response of different weed management practices to the growth and yield of black gram. *Cyperus* sp., *Phyllanthus niruri, Trianthema portulacastrum, Amaranthus viridis* were the dominant weeds. It was observed that besides weed-free treatment the lowest weed count, dry weight of weeds (3.66 g m⁻²), weed index (2.73%) as well as highest weed control efficiency (60.06%) were found with the pre-emergence (PE) application of Pendimethalin 1 kg ha⁻¹ followed by one hand weeding at 40 days after sowing (DAS). The maximum growth attributes, yield attributes and yield were found with the pre-emergence application of Pendimethalin 1 kg a.i. ha⁻¹ followed by one hand weeding at 40 DAS which was statistically at par with pre-emergence application of Pendimethalin 1 kg a.i. ha⁻¹ followed by one hand weeding at 40 days after sowing explication of imazethapyr 75 g a.i. ha⁻¹ at 40 days after sowing. The economic parameters were recorded highest with the application of Pendimethalin (PE) 1 kg a.i. ha⁻¹ followed by one-hand weeding at 40 days after sowing. Hence Pendimethalin is the most effective treatment among the other treatments.

Keywords: Blackgram, chemical control, integrated weed management, pendimethalin, growth, yield

Introduction

Black gram (*Vigna mungo* L.) is one of the important pulses grown in different agro-ecological areas throughout the country in the summer and rainy seasons. It is also called Urd bean or Mash bean and belongs to the family Leguminaceae and subfamily Papilionaceae. It is the fourth most important pulse crop in India after chickpeas, pigeon pea and green gram. It is widely cultivated in terms of area in Rajasthan (38.39 lakh ha), Madhya Pradesh (21.14 lakh ha), Maharashtra (19.28 lakh ha), Karnataka (18.32 lakh ha), Uttar Pradesh (8.53 lakh ha), Jharkhand (4.23 lakh ha), Tamil Nadu (1.79 lakh ha) In Punjab, it was grown on 2.0 thousand hectare in 2019-20 and total production was recorded 1.2 thousand tons (Anonymous. 2020) ^[1].

Black gram has a wide range of adaptability and the ability to withstand stress conditions. It is a self-pollinated legume crop. It contains 24% protein, 60% carbohydrates, 1.3% fat, 3.2% minerals, 0.9% fiber, 154 mg calcium, 385 mg phosphorus, 9.1 mg iron and a small amount of vitamin B complex (Anonymous 2020)^[1]. It can increase soil fertility because it can fix around 70-90 kg N ha⁻¹. After removing the pods, the plants can be used as high-quality green fodder or dry fodder or green manure (Carvalho *et al.* 2015)^[3].

There are various factors that contribute to the low yield of black gram. Among those various factors, weeds are considered the most important. The losses caused by weeds exceed the losses caused by any other type of agricultural pests, such as insects, nematode, disease, rodents etc. The crop is not a good competitor against weeds during the early stage (Choudhary *et al.* 2012)^[2]. Black gram is generally grown in rainfed conditions during the *kharif* season, where weeds are the main cause of low crop productivity. Weeds at critical periods of crop-weed competition caused reduction of 80-90% in yield depending upon type and intensity of weed infestation (Kumar *et al.* 2018)^[6]. Uncontrolled weeds have been reported to cause a considerable reduction (46-53%) in seed yield of black gram (Singh *et al.* 2016)^[9].

To control weeds, the traditional method of weed control i.e. hand weeding although is very effective but it is expensive, tedious and time-consuming (Yadav *et al.* 2009) ^[11]. Due to this weeds are controlled by mechanical, chemical and biological methods which can be used alone or in combination with more than one method. The chemical method of weed control is not only cost effective but also efficient in minimizing weed infestation for longer period provided they are applied judiciously. Imazethapyr, a broad-spectrum herbicide, has soil and foliar activity that allows flexibility in its application timing and has low mammalian toxicity (Tan *et al.* 2005) ^[10]. Pendimethalin is basically a pre-emergence herbicide that can effectively control weeds in beans.

Materials and Methods

The field experiment was conducted during the summer season of the year 2021-22 at Campus for Research and Advanced Studies, Dhablan, G.S.S.D.G.S. Khalsa College, Patiala. The field experiment is situated at about 30°19' North latitude, 76° 24' East longitude and altitude of 247 meter above the mean sea level. The cultivar of Black gram MASH 1008 was sown in lines

during the second week of April at a spacing of 30×10 cm with the help of a hand seed drill following all the recommended packages and practices (Anonymous 2020)^[1]. The soil of the experimental field was clayey in texture having pH 7.3. The soil was medium in available nitrogen (295.13 kg ha⁻¹), phosphorous (18.25 kg ha⁻¹) and potassium (157.59 kg ha⁻¹). The organic carbon was (0.6%). There were 11 weed control treatments viz. weedy check, weed free, one hand weeding (HW) at 20 days after sowing (DAS), two HW at 20 and 40 DAS, Pendimethalin (PE) 1 kg ha⁻¹, Pendimethalin (PE) 1kg ha⁻¹ followed by one HW at 40 DAS, imazethapyr (PoE) 0.075 kg ha⁻¹ at 20 DAS. quizalofop-ethyl (PoE) 0.050 kg ha-1 at 20 DAS. Pendimethalin (PE) 1kg ha⁻¹ followed by imazethapyr (PoE) 0.075 kg ha⁻¹ at 40 DAS, quizalofop-ethyl (PoE) 0.050 kg ha⁻¹at 20 DAS + HW at 40 DAS, Straw mulch (5 t ha⁻¹) and making the 33 experimental units. The data regarding weed and growth parameters were recorded at 25 and 50 DAS and at the harvest stage. The data on number of weeds and dry weight of weeds were transformed using $(\sqrt{X+0.5})$ before subjected to statistical analysis and then weed control efficiency (WCE) was calculated. The experiment was laid out in randomized block design with three replications.

Details of layout

Experimental design	Randomized Block Design
No. of replications	3
No. of treatments	11
Total number of plots	33
Spacing	$30 \times 10 \text{ cm}$
Gross plot size	$3.6 \times 4.05 \text{ m}$
Net plot size	3.3 × 3.9 m
Variety	MASH 1008

T1	Weedy check
T ₂	Weed free
T 3	One HW at 20 DAS
T_4	Two HW at 20 and 40 DAS
T ₅	Pendimethalin (PE) 1 kg ha ⁻¹
T ₆	Pendimethalin (PE) 1 kg ha ⁻¹ fb one HW at 40 DAS
T 7	Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 20 DAS
T8	Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS
T9	Pendimethalin (PE) 1 kg ha ⁻¹ fb Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 40 DAS
T ₁₀	Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS + HW at 40 DAS
T ₁₁	Straw mulch (5 t ha ⁻¹)

Treatment Details

Results and Discussion

Effect of different weed management practices on weeds population, dry weight of weeds (g m⁻²), weed control efficiency (%) and weed index (%) at harvest of black gram. The predominant weeds noticed in blackgram field were Cyperus sp., *Phyllanthus niruri, Trianthema portulacastrum, Amaranthus viridis.* As a result of this experiment, it was observed that beside weed free treatment, the lowest number of weeds, dry weight of weeds (3.66 g m⁻²) and weed index (2.73%), highest weed control efficiency (60.06%) was recorded with an application of Pendimethalin (PE) 1 kg ha⁻¹ followed by one HW at 40 DAS as shown in table no. 2. This might be due to the initial flush of weed germination being restricted by Pendimethalin herbicide and in later stage weeds were controlled by manual weeding. Our results are in comparison with the study conducted by Kumar *et al.* (2005) ^[12] in which it was recorded that with the application of Pendimethalin (PE) in combination with raised seed bed planting significantly less number of weeds along with increased yield of black gram were obtained.

Table 1: Effect of different	weed management	practices on wee	eds population at 1	harvest of black gram
	0	1	1 1	0

Treatments		Phyllanthus niruri	Trianthema portulacastrum	Amaranthus viridis
T ₁ : Weedy check	4.81 (22.17)	3.89 (14.17)	4.25 (17.10)	4.43 (18.68)
T ₂ : Weed free	0	0	0	0
T ₃ : One HW at 20 DAS	4.20 (16.67)	2.97(7.85)	3.5(11.25)	3.95 (14.65)
T4: Two HW at 20 and 40 DAS	3.76 (13.18)	2.66 (6.10)	2.99 (7.95)	3.06 (8.41)
T ₅ : Pendimethalin (PE) 1 kg ha ⁻¹	4.27 (17.28)	3.02 (8.16)	3.59(11.93)	3.73 (12.95)
T ₆ : Pendimethalin (PE) 1 kg ha ⁻¹ fb one HW at 40 DAS	3.45 (10.95)	2.42 (4.87)	2.71 (6.35)	2.93 (7.62)
T ₇ : Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 20 DAS	3.93 (14.52)	2.78(6.73)	3.22 (9.37)	3.34 (10.16)
T ₈ : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS	4.01 (15.16)	2.83 (7.05)	3.3 (9.89)	3.37 (10.36)
T ₉ : Pendimethalin (PE) 1 kg ha ⁻¹ <i>fb</i> Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 40 DAS	3.65 (12.37)	2.57(5.65)	2.83(7.03)	3.02 (8.14)
T10: Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS + HW at 40 DAS	3.86 (13.91)	2.73 (6.47)	3.10 (8.64)	3.20 (9.26)
T_{11} : Straw mulch (5 t ha ⁻¹)	4.11 (15.95)	2.91 (7.52)	3.38 (10.47)	3.45 (10.94)
S.E(d)±	0.11	0.23	0.12	0.5
C.D. at 5%	0.26	0.56	0.25	0.12

 Table 2: Effect of different weed management practices on dry weight of weeds (g m-2), weed control efficiency (%), weed index (%) at harvest of black gram

Treatments	Dry weight of weeds (g m- ²)	Weed control efficiency (%)	Weed index (%)
T ₁ : Weedy check	7.03 (48.46)	0	44.26
T ₂ : Weed free	0	100.00	0
T ₃ : One HW at 20 DAS	5.18 (25.85)	32.88	24.65
T ₄ : Two HW at 20 and 40 DAS	4.01 (15.10)	52.55	15.94
T ₅ : Pendimethalin (PE) 1 kg ha ⁻¹	5.45 (28.75)	33.01	35.45
T ₆ : Pendimethalin (PE) 1 kg ha ⁻¹ fb one HW at 40 DAS	3.66 (12.41)	60.06	2.73
T ₇ : Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 20 DAS	4.52 (19.42)	45.71	19.96
T ₈ : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS	4.63 (20.51)	43.47	22.14
T9: Pendimethalin (PE) 1 kg ha ⁻¹ fb Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 40 DAS	3.96 (14.75)	56.81	11.31
T ₁₀ : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS + HW at 40 DAS	4.2 (16.64)	48.71	19.36
T_{11} : Straw mulch (5 t ha ⁻¹)	4.77 (21.89)	40.25	31.51
$S.E(d)\pm$	0.14	1.68	0.08
C.D. at 5%	0.32	3.50	0.16

Effect of different weed management practices on growth parameters of black gram

Besides weed-free treatment, maximum plant height (39.50 cm), dry weight plant⁻¹ (11.94 g), number of branches plant⁻¹ (6.48), leaf area index (1.31) were recorded with an application of Pendimethalin (PE) 1 kg ha⁻¹ followed by one HW at 40 DAS. The minimum values of all the growth parameters were observed under the weedy check. There is lesser crop-weed competition which resulted in the higher utilization of available nutrients and moisture by the crop that results in the vigorous growth of black gram plants. Khot *et al.* (2016) ^[8] and Komal *et al.* (2015) ^[7] conducted an experiment to study the effect of different sowing time and weed management practices with respect to yield, quality and nutrient uptake on summer black gram and observed significant reduction in weed growth with the application of Pendimethalin followed by one HW at 40 DAS.

Table 3: Effect of different weed management practices on growth parameters of black gram

Traatmants		Dry weight	Number of	Leaf area
Treatments	(cm)	plant ⁻¹	branches plant ⁻¹	index
T ₁ : Weedy check	24.73	7.20	4.83	0.97
T ₂ : Weed free	40.90	12.17	7.10	1.43
T ₃ : One HW at 20 DAS	26.53	8.55	5.42	1.04
T ₄ : Two HW at 20 and 40 DAS	35.10	10.31	6.16	1.25
T ₅ : Pendimethalin (PE) 1 kg ha ⁻¹	25.57	7.80	5.25	0.99
T ₆ : Pendimethalin (PE) 1 kg ha ⁻¹ followed by one HW at 40 DAS	39.50	11.94	6.48	1.31
T ₇ : Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 20 DAS	31.27	9.53	5.67	1.18
T ₈ : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS	30.23	9.08	5.59	1.15
T ₉ : Pendimethalin (PE) 1 kg ha ⁻¹ followed by Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 40 DAS	37.87	10.90	6.25	1.29
T_{10} : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS + HW at 40 DAS	33.27	9.93	5.98	1.20
T_{11} : Straw mulch (5 t ha ⁻¹)	28.27	8.84	5.48	1.09
SE (d)±	2.15	0.79	0.16	0.02
C.D. at 5%	4.48	1.65	0.33	0.03

Effect of different weed management practices on the yield parameters of black gram

The yield parameters were significantly influenced with an

application of different weed control parameters. The maximum yield attributing parameters such as number of pods $plant^{-1}$ (25.22), number of seed pod^{-1} (5.69) and seed index (4.72 g) as

well as grain yield (12.81 q ha⁻¹), straw yield (27.84 q ha⁻¹), biological yield (38.65 q ha⁻¹) and harvest index (33.06%) were recorded with the application of Pendimethalin (PE) 1 kg ha⁻¹ followed by one HW at 40 DAS. While the minimum yield was recorded from the weedy check. It might be due to the effective integrated weed control treatments as it minimized the state of crop-weed competition, so that the plants do not face the moisture or nutrients stress conditions that might be due to lesser weed infestation. With a better nutrient supply and weed competition free optimum soil environment, the seed size of crop ultimately improves and results in the enhancement of all the yield parameters.

Table 4: Effect of different weed manageme	ent practices on the	yield parameters o	of black gram
--	----------------------	--------------------	---------------

Treatments	No. of	No. of	Seed	Grain yield	Straw yield	Biological	Harvest
Treatments	pods plant ⁻¹	seed pod ⁻¹	index (g)	(q ha ⁻¹)	(q ha ⁻¹)	yield (q ha ¹)	index (%)
T ₁ : Weedy check	21.64	4.97	4.07	7.34	18.85	26.19	28.06
T ₂ : Weed free	25.40	5.80	4.80	13.17	29.22	42.39	31.07
T ₃ : One HW at 20 DAS	23.48	5.15	4.29	9.91	23.49	33.40	29.69
T4: Two HW at 20 and 40 DAS	24.96	5.53	4.63	11.07	25.25	36.32	30.49
T ₅ : Pendimethalin (PE) 1 kg ha ⁻¹	23.02	5.09	4.20	8.50	21.08	29.58	28.76
T ₆ : Pendimethalin (PE) 1 kg ha ⁻¹ followed by one HW at 40 DAS	25.22	5.69	4.72	12.81	27.84	38.65	33.06
T ₇ : Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 20 DAS	24.43	5.38	4.51	10.54	24.46	35.00	30.10
T ₈ : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS	24.03	5.29	4.44	10.24	24.21	34.45	29.71
T9: Pendimethalin (PE) 1 kg ha ⁻¹ followed by Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 40 DAS	25.08	5.60	4.67	11.68	26.49	37.28	31.34
T ₁₀ : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS HW at 40 DAS	24.81	5.43	4.56	10.62	24.87	35.49	29.93
T ₁₁ : Straw mulch (5 t ha ⁻¹)	23.70	5.22	4.37	9.02	21.91	30.93	29.11
SE (d)±	0.13	0.098	0.29	0.55	0.72	0.75	0.86
C.D. at 5%	0.26	0.20	0.62	1.15	1.51	1.57	1.80

Effect of different weed management practices on economy of black gram

The economic parameters such as gross return, net return and benefit cost ratio (B:C ratio) was significantly influenced by all

the weed treatments. The highest gross return (Rs. 80703), net return (Rs.55297) and B:C ratio (2.18) was observed with an application of Pendimethalin (PE) 1 kg ha⁻¹ followed by one HW at 40 DAS.

Table 5: Effect of different weed management practices on economy of black gram

Treatments	Gross return (Rs. ha ⁻¹)	Net return (Rs. ha ⁻¹)	B:C ratio
T ₁ : Weedy check	46,242	25,392	1.21
T ₂ : Weed free	82,971	53,721	1.84
T ₃ : One HW at 20 DAS	62,433	38,783	1.64
T4: Two HW at 20 and 40 DAS	69,741	43,291	1.64
T ₅ : Pendimethalin (PE) 1 kg ha ⁻¹	53,550	30,944	1.37
T ₆ : Pendimethalin (PE) 1 kg ha ⁻¹ followed by one HW at 40 DAS	80,703	55,297	2.18
T ₇ : Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 20 DAS	66,402	44,542	2.04
T ₈ : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS	64,512	41,502	1.80
T ₉ : Pendimethalin (PE) 1 kg ha ⁻¹ followed by Imazethapyr (PoE) 0.075 kg ha ⁻¹ at 40 DAS	73,584	49,978	2.12
T ₁₀ : Quizalofop-ethyl (PoE) 0.050 kg ha ⁻¹ at 20 DAS + HW at 40 DAS	66,906	41,096	1.59
T_{11} : Straw mulch (5 t ha ⁻¹)	56,826	32,076	1.30
SE (d)±	3464	3464	0.13824
C.D. at 5%	7226	7226	0.28836

Conclusion

The current study revealed that among all the treatments, the application of Pendimethalin (PE) 1 kg ha⁻¹ followed by one HW at 40 DAS gives the profitable results with respect to the growth, yield as well as economic parameters and was found most efficient treatment to control the weeds.

References

- Anonymous. Package of Practices for Kharif Crops of Punjab. Punjab Agricultural University Ludhiana; c2020. p. 80-85.
- 2. Choudhary, Kumar VK, Bhagawati R. Integrated weed management in blackgram (*Vigna mungo*) under mid hills of Arunachal Pradesh. Indian Journal Agronomy. 2012;57(4):382-385.
- 3. Carvalho NS, Oliveira ABB, Pessoa MMC, Costa Neto VP,

De Sousa RS, Cunha JR, *et al.* Short-term effect of different green manure on soil chemical and biological properties, African Journal of Agricultural Research. 2015;10:4076-4081.

- 4. Devaraju B, Senthivel T. Weed Management in irrigated Black gram (*Vigna mungo* L) in southern zone of Tamil Nadu. Journal of Advance Studies in Agricultural, Biological and Environmental Sciences. 2017;4(4):29-33.
- Chandravanshi MD, Patel JR, Kasyap S. Effect of integrated nutrient management on growth and yield of Urdbean (*Vigna mungo* L.) in Chhattisgarh plain. Int. J Adv. Chem. Res. 2022;4(2):260-265. DOI: 10.33545/26646781.2022.v4.i2d.109
- 6. Kumar S, Mishra P, Singh P, Sinha K. Effect of different weed control practices on growth of Urd Bean. International Journal of Chemical Studies. 2018;6(1):1857-1862.

- 7. Komal, Singh SP, Yadav RS. Effect of weed management on growth, yield and nutrient uptake of green gram. Indian Journal of Weed Sciences. 2015;42(7):206-210.
- 8. Khot AB, Sagvekar VV, Muthal YC, Panchal VV, Dhonde MB. Effect on summer black gram (*Phaseolus mungo* L.) to different sowing time and weed management practices with respect to yield, quality and nutrient uptake. International Journal of Tropical Agriculture. 2016;34(7):2155-2161.
- Singh VP, Singh TP, Singh SP, Kumar A, Satyawali K, Banga A, *et al.* Weed management in blackgram with premix herbicides. Indian Journal of Weed Sciences. 2016;48(2):178-181.
- Tan S, Evans RR, Dahmer ML, Singh BK, Shaner DL. Imidazolin one-tolerant crops: history, current status and future. Pest Management Sciences. 2005;61(2):246-257.
- 11. Yadav MK, Singh RS. Effect of nitrogen levels and weed management practices on pigeon pea (*Cajanus cajan*) and rice (*Oryza sativa*) intercropping system under ridge-furrow planting system. Indian Journal Agricultural Sciences. 2009;79(4):268-272.
- Kumar S, Angiras N. Efficacy of Pendimethalin under different planting methods of blackgram (*Vigna mungo*). Indian Journal of Weed Sciences. 2005;37(3-4):216-219.