



# International Journal of Research in Agronomy

E-ISSN: 2618-0618

P-ISSN: 2618-060X

© Agronomy

[www.agronomyjournals.com](http://www.agronomyjournals.com)

2023; 6(1): 26-30

Received: 13-10-2022

Accepted: 19-11-2022

**Roshan Kumawat**

Ph.D., Scholar, Department of  
Agronomy, Agriculture University,  
Borkheda, Kota, Rajasthan, India

**AC Shivran**

Professor and O.S. D., SKNAU,  
Jobner, Rajasthan, India

**Rajesh Kumar**

Assistant Professor, Department of  
Agronomy, Agriculture University,  
Borkheda, Kota, Rajasthan, India

**Anju Bijarnia**

Ph.D., Scholar, Department of  
Agronomy, Agriculture University,  
Borkheda, Kota, Rajasthan, India

**Lali Jat**

Ph.D., Scholar, Department of  
Agronomy, SPVUAT, Meerut,  
Uttar Pradesh, India

## Yield and quality of mustard (*Brassica juncea*) and lentil (*Lens esculenta*) intercropping system influenced by different fertility levels

**Roshan Kumawat, AC Shivran, Rajesh Kumar, Anju Bijarnia and Lali Jat**

**DOI:** <https://doi.org/10.33545/2618060X.2023.v6.i1a.163>

### Abstract

A field experiment was conducted in Split Plot Design with five intercropping systems [sole mustard, sole lentil, mustard + lentil (1:3), mustard + lentil (2:2) and mustard + lentil (2:3)] and four fertility levels (0, 75, 100 and 125% RDF) and replicated thrice during *rabi* season, 2017-18 at Agronomy farm, S.K.N. College of Agriculture, Jobner. The results revealed that planting of mustard in 2:2 row ratio, being at par to sole mustard, gave significantly higher seed yield (2234 kg/ha) as compared to 1:3 and 2:3 row ratios. The 1:3 row ratio, remaining at par with sole lentil, recorded significantly higher seed yield of lentil as compared to 2:2 and 2:3 row ratio. The 1:3 row ratio, recorded significantly higher nitrogen concentration in seed of mustard. Planting of mustard in 2:2 and planting of lentil in 1:3 row ratio, recorded significantly highest total uptake of nitrogen and phosphorus. Application of fertility levels up to 125% RDF significantly increased the seed yield of mustard and lentil and nitrogen concentration in the seed of mustard and straw of lentil, total nitrogen and phosphorus uptake of mustard and lentil. Whereas, RDF up to 100% significantly increased nitrogen concentration in stover and seed of lentil, phosphorus concentration in seed and stover/straw of mustard and lentil, oil content in seed of mustard and protein content in seed of lentil.

**Keywords:** Mustard, Lentil, Intercropping, RDF, Protein, Oil

### 1. Introduction

Oilseed crops are the main source of energy in the diet of Indians. Though, India has become self-reliant with respect to food grains but still lagging behind in the production of oilseeds. Indian mustard (*Brassica juncea*) is one of the major edible oil seed crop, maximizing mustard production and productivity is crucial for reducing the edible oil import burden for the country. While lentil is a good supplier of protein and has a high nutritional value, its production and productivity enhancement is vital for achieving food and nutritional security. Hence, maximizing the productivity of Indian mustard (*Brassica juncea*) + lentil (*Lens esculenta*) cropping systems is crucial for the country. Intercropping is one of the most important techniques which, involves the growing of crops under different plant geometry. To avoid the risk of sole crop failure, adoption of intercropping is more safe and profitable cropping system for increasing the total production and net profit per unit area. Intercropping offers to farmers an opportunity to utilize diverse resources available at their farm (Ghosh, 2004) [7]. Legumes offer excellent compatible combinations for mixing with oilseeds to minimize competition and to confer symbiotic association to achieve the prime aim of maximum use of available resources. The productivity of crop is largely influenced by fertility management. The dose of fertilizer depends on the initial soil fertility status and moisture availability conditions. Exploitation of genetic yield potentiality of new plant types of mustard although depends on the extent of fertilizer application, the balance and compatible composition of all the major nutrients is a must to obtain greater fertilizer use efficiency. Among the commonly applied major nutrients, nitrogen is the key element in mustard production, which is a structural components of protein molecules, amino acids, chlorophyll and other constituents. Its adequate supply promotes higher photosynthesis activity and vigorous vegetative growth. A higher nitrogen supply favours the conversion of carbohydrates into protein.

**Corresponding Author:**

**Roshan Kumawat**

Ph.D., Scholar, Department of  
Agronomy, Agriculture University,  
Borkheda, Kota, Rajasthan, India

Phosphorus plays an important role in improving water use efficiency due to its favourable effect on crop growth root development and high seed yield production per unit of water use. Its conspicuous role in hastening crop maturity is of still more interest to us in the age of intensive cropping systems.

## 2. Materials and Methods

The experiment was conducted at the Agronomy farm (Plot No. 1-C) S.K.N. College of Agriculture, Jobner (Rajasthan) during *rabi* season 2017-18. The experiment comprising twenty treatment combinations replicated thrice in Split Plot Design (SPD) with five intercropping systems [sole mustard, sole lentil, mustard + lentil (1:3), mustard + lentil (2:2) and mustard + lentil (2:3)] in main plot and four fertility levels (0, 75, 100 and 125% RDF) in sub plot. The soil of experimental field was loamy sand in texture, alkaline in reaction (pH 8.2), poor in organic carbon (0.14%), low in available nitrogen (130.0 kg/ha) and medium in phosphorus (16.52 kg P<sub>2</sub>O<sub>5</sub>/ha) and potassium (151.8 kg/ha) content. The seeds of mustard variety T-59 @ 5 kg/ha and Lentil variety RLG-255 @ 40 kg/ha were used for sowing in the experiment. The sowing was done on 09.11.2017. Mustard seeds were sown at a spacing of 45 x 15 cm<sup>2</sup> and lentil was spaced at an intra-row spacing of 10 cm with 30 cm inter-row spacing. In 1:3 row ratio, three rows of lentil were planted after replacing one row of mustard in the replacement series. In 2:2 and 2:3 row ratios, two and three rows of lentil were planted in between paired rows of mustard as additive series. Half dose of nitrogen through urea and full dose of phosphorus through DAP were incorporated in the soil before sowing in plots as per treatments and the remaining half dose of nitrogen was applied 40 DAS. 60 kg N and 30 kg P<sub>2</sub>O<sub>5</sub>/ha for mustard sole and 20 kg N and 40 kg P<sub>2</sub>O<sub>5</sub>/ha for sole crop of lentil were applied. In intercropping systems, the RDF of mustard was applied. Yield of mustard and lentil were computed from the plants of net plot in each treatment. Oil percent in the mustard was determined by Soxhlet apparatus using petroleum ether (60-80 °C) as an extractant and protein content in seed of lentil crop was calculated by multiplying the per cent N content in seed with the factor of 6.25 (A.O.A.C., 1960) [1].

The uptake of nitrogen by crop was calculated by following formula (Snell and Snell, 1949) [16]:

$$N \text{ uptake (kg/ha)} = \frac{\text{Per cent N in seed} \times \text{Seed yield (kg/ha)} + \text{Per cent N in stover/straw} \times \text{stover/straw yield (kg/ha)}}{100}$$

The uptake of phosphorus by crop was calculated by following formula (Jackson, 1967) [8]:

$$P \text{ uptake (kg/ha)} = \frac{\text{Per cent P in seed} \times \text{Seed yield (kg/ha)} + \text{Per cent P in stover/straw} \times \text{stover/straw yield (kg/ha)}}{100}$$

$$\text{Total uptake (kg/ha)} = \frac{\text{Nutrient conc. in seed (\%)} \times \text{Seed yield (kg/ha)} + \text{Nutrient conc. in stover/straw (\%)} \times \text{stover/straw yield (kg/ha)}}{100}$$

## 3. Results and discussion

### 3.1 Effect of intercropping

#### 3.1.1 Yield

Results indicated that planting of mustard in 2:2 row ratio with lentil obtained significantly the highest seed yield (2234 kg/ha) of mustard as compared to 1:3 and 2:3 row ratios, whereas it remained statistically at par with sole planting of mustard. The

2:2 row ratio also improved the seed yield of mustard by 77.2 and 16.1 per cent over 1:3 and 2:3 row ratios, respectively. It may be due to 100% population and beneficial effect of lentil as evidenced by increased growth and yield attributes. These findings are in close conformity with those of Meena *et al.* (2008) [11], Singh *et al.* (2014) [15] and Das *et al.* (2017) [5].

A significant reduction in seed yield of lentil was recorded in 2:2 (739 kg/ha) and 2:3 (980 kg/ha) row ratios as compared to 1:3 (1441 kg/ha) row ratio and 1:3 row ratio remained at par with sole lentil (1400 kg/ha). The reduction in seed yield of lentil in 2:2 and 2:3 row ratios over sole lentil was primarily due to low plant population of lentil in 2:2 row ratio and a significant reduction in yield attributes (pods/plant and seeds/pod). This might also be due to adverse effect of mustard as mustard was planted in paired row with 100% population and lentil was added between pair as additive series. Similar results were also found by Yadav *et al.* (2014) [18] and Das *et al.* (2017) [5].

#### 3.1.2 Nutrient concentration, uptake and quality

A significant increase in nitrogen concentration in seed of mustard was recorded in 1:3 row ratio as compared to 2:3 row ratio, sole mustard and 2:2 row ratio. This was probably due to the comparatively greater availability of nutrients to mustard in 1:3 row ratio as compared to 2:3 row ratio, sole mustard and 2:2 row ratio. This might be due to the complementary effect of intercropping. The results are in close conformity with those of Devi *et al.* (2014) [16]. Among different intercropping systems, the planting of mustard in 2:2 row ratio gave significantly higher total nitrogen and phosphorus uptake as compared to 1:3 and 2:3 row ratio as well as sole crop. Further, sole mustard and 2:3 row ratio also gave significantly higher total nitrogen and phosphorus uptake as compared to 1:3 row ratio.

Similarly, planting of lentil in 1:3 row ratio also gave significantly higher total nitrogen and phosphorus uptake by lentil crop as compared to rest of intercropping treatments and remained at par with sole crop of lentil. Because nutrient uptake is a function of its concentration in crop plant and seed and stover/straw yields of the crop. The results are in close conformity with those of Meena *et al.* (2008) [11] and Abraham and Thenua (2015) [2]. Oil content in the seed of mustard and protein content in seed of lentil were not improved significantly by different systems of intercropping.

### 3.2 Effect of fertility levels

#### 3.2.1 Yield

Application of Increasing levels of fertility up to 125% RDF recorded significant improvement in seed (2129 kg/ha of mustard and 1309 kg/ha of lentil), stover/straw (6428 kg/ha of mustard and 3663 kg/ha of lentil) and biological (8557 kg/ha of mustard and 4972 kg/ha of lentil) yields of mustard and lentil. The increase in seed and stover/straw yields might be due to better nutritional status in the soil, which was low in N and P. The increased supply of NPK and their higher uptake by plants might have stimulated the rate of various physiological processes in plants and led to increased growth and yield parameters and ultimately resulted in higher seed and stover/straw yields. The biological yield is a function of seed and stalk yields. Thus, a significant increase in biological yield with the application of N and P could be ascribed due to increased seed and stover/stalk yields. The results of the present investigation are in line with those of Meena *et al.* (2013) [10], Sharma *et al.* (2013) [14], Singh *et al.* (2014) [15], Balai *et al.* (2017b) [4], and Keerthi *et al.* (2017) [9].

### 3.2.2 Nutrient concentration, uptake and quality

The significant increase in N concentration in the seed of mustard and straw of lentil and total nitrogen and phosphorus uptake of mustard and lentil were recorded with increased application of RDF up to 125%. The nitrogen concentration in stover of mustard and in seed of lentil, phosphorus concentration in seed and stover/straw, oil content in seed of mustard and protein content in seed of lentil were recorded as significantly higher with increased application of 100% RDF over lower levels. This might be due to an improved nutritional environment in the rhizosphere as well as in the plant system leading to enhanced translocation of N and P in plant parts. Since the nutrient uptake is a function of its content in crop plant

and seed and stover/straw yields of the crop. The increase in these parameters due to N and P fertilization led to increased uptake of nutrients in the present study. Significant increase in protein content and oil percentage because of increased N content in seed which might be the result of increased availability of nitrogen to plants. Another reason for higher nitrogen content might be due to the increased activity of the nitrate reductase enzyme. Higher nitrogen in seed is directly responsible for higher protein because it is a primary component of amino acids that constitute the basis of protein. These results are in close conformity with the findings of Parihar *et al.* (2009)<sup>[13]</sup>, Trivedi *et al.* (2013)<sup>[17]</sup>, Pandey *et al.* (2016)<sup>[12]</sup> and Balai *et al.* (2017a)<sup>[3]</sup>.


**Table 1:** Effect of the intercropping system and fertility levels on yields, nitrogen and Phosphorus concentration in seed and stover and total nitrogen and Phosphorus uptake and oil content in the seed of mustard

Treatment	Yields (kg/ha)		Nitrogen Concentration (%)		Total nitrogen uptake (kg/ha)	Oil content (%)	Phosphorus concentration (%)		Total phosphorus uptake (kg/ha)
	Seed	Stover	Seed	Stover			Seed	Stover	
A. Intercropping system									
Mustard sole	2050	6214	3.18	0.748	113.01	35.33	0.660	0.256	29.76
Mustard + Lentil (1:3)	1261	3776	3.91	0.787	79.99	37.29	0.700	0.268	19.15
Mustard + Lentil (2:2)	2234	6823	3.58	0.772	134.25	36.64	0.692	0.262	33.70
Mustard + Lentil (2:3)	1924	5795	3.09	0.733	103.06	35.05	0.648	0.253	27.43
S.Em±	53	177	0.09	0.019	2.61	0.88	0.017	0.006	0.65
CD (P=0.05)	185	612	0.31	NS	9.02	NS	NS	NS	2.26
B. Fertility level									
Control	1557	4688	2.90	0.648	74.92	32.92	0.573	0.223	19.32
75% RDF	1800	5458	3.33	0.742	99.62	35.73	0.657	0.254	25.62
100% RDF	1983	6035	3.64	0.811	120.14	37.82	0.722	0.278	31.01
125% RDF	2129	6428	3.89	0.839	135.62	37.85	0.748	0.284	34.09
S.Em±	36	102	0.06	0.014	2.07	0.60	0.013	0.005	0.54
CD (P=0.05)	118	334	0.21	0.047	6.76	1.97	0.042	0.016	1.75


NS= Non-significant

**Table 2:** Effect of intercropping system and fertility levels on yields, nitrogen and Phosphorus concentration in seed and straw and total nitrogen and Phosphorus uptake and protein content in seed of lentil

Treatment	Yields (kg/ha)		Nitrogen Concentration (%)		Total nitrogen uptake (kg/ha)	Protein content (%)	Phosphorus concentration (%)		Total phosphorus uptake (kg/ha)
	Seed	Straw	Seed	Straw			Seed	Straw	
A. Intercropping system									
Lentil sole	1400	3901	3.51	1.59	112.41	21.94	0.342	0.214	13.31
Mustard + Lentil (1:3)	1441	4021	3.57	1.62	117.89	22.31	0.345	0.218	13.92
Mustard + Lentil (2:2)	739	2060	3.42	1.53	57.43	21.38	0.328	0.209	6.82
Mustard + Lentil (2:3)	980	2730	3.37	1.49	74.53	21.06	0.321	0.203	8.80
S.Em±	33	97	0.08	0.04	2.84	0.52	0.008	0.005	0.34
CD (P=0.05)	113	335	NS	NS	9.82	NS	NS	NS	1.18
B. Fertility level									
Control	937	2604	3.11	1.31	63.63	19.44	0.281	0.176	7.26
75% RDF	1097	3054	3.41	1.50	83.71	21.31	0.324	0.205	9.87
100% RDF	1217	3391	3.66	1.65	101.10	22.88	0.357	0.226	12.08
125% RDF	1309	3663	3.69	1.77	113.82	23.06	0.374	0.237	13.65
S.Em±	23	68	0.07	0.03	1.91	0.43	0.007	0.004	0.23
CD (P=0.05)	76	221	0.23	0.10	6.24	1.41	0.022	0.014	0.74



**Fig 1:** Effect of intercropping system and fertility levels on total nitrogen and phosphorus uptake and oil content in seed of mustard



**Fig 2:** Effect of intercropping system and fertility levels on total nitrogen and phosphorus uptake and protein content in seed of lentil

#### 4. References

1. AOAC. Official method of analysis, 18 edition. Association of Official Agricultural Chemists, Washington; c1960.
2. Abraham T, Thenua OVS. Impact assessment of different irrigation and fertilizer levels on mustard + chickpea intercropping systems over sole cropping. International Journal of Modern Chemistry and Applied Sciences. 2015;2(4):266-272.
3. Balai K, Jajoria M, Verma R, Deewan P, Bairwa SK. Nutrient content, uptake, quality of chickpea and fertility status of soil as influenced by fertilization of phosphorus and zinc. Journal of Pharmacognosy and Phytochemistry. 2017a;6(1):392-398.
4. Balai K, Sharma Y, Jajoria M, Deewan P, Verma R. Effect of phosphorus and zinc on growth, yield and economics of chickpea (*Cicer arietinum* L.). International Journal of Current Microbiology and Applied Sciences. 2017b;6(1):1174-1181.
5. Das S, Devi KN, Athokpam HS, Lhungdim J, Longjam M. Chickpea (*Cicer arietinum* L.) based intercropping system with rapeseed (*Brassica juncea* L.) on growth, yield and competition indices. Environment and Ecology. 2017;35(1B):470-430.
6. Devi KN, Shamurailatpam D, Singh TB, Athokpam HS, Sing NB, Singh NG, et al. Performance of lentil (*Lens culinaris* M.) and mustard (*Brassica juncea* L.) intercropping under rainfed conditions. Australian Journal of Crop Science. 2014;8(2):284-289.
7. Ghosh PK. Growth, yield, competition and economics of groundnut/cereal fodder intercropping systems in the semi-arid tropics of India. Field Crops Research. 2014;88(2-3):227-237.
8. Jackson ML. Soil chemical analysis. Prentice Hall Inc. N.J. USA; c1967.
9. Keerthi P, Pannu RK, Dhaka AK, Daniel J, Yogesh. Yield, nitrogen uptake and nutrient use efficiency in Indian mustard (*Brassica juncea* L.) as effected by date of sowing and nitrogen levels in western Haryana, India. International Journal of Current Microbiology and Applied Sciences. 2017;6(4):1168-1177.
10. Meena DS, Meena VR, Meena AK. Fertilizer management studies on growth and productivity of hybrid Indian mustard (*Brassica juncea* L.). Journal of Oilseed Brassica. 2013;4(1):39-42.
11. Meena SL, Shamsudheen M, Dayal D. Impact of row ratio and nutrient management on performance of clusterbean (*Cyamopsis tetragonoloba*) + sesame (*Sesamum indicum*) intercropping system. Indian Journal of Agronomy. 2008;53(4):285-289.
12. Pandey MK, Verma A, Kumar P. Effect of integrated phosphorus management on growth, yield and quality of lentil (*Lens culinaris*). Indian Journal of Agricultural Research. 2016;50(3):238-243.
13. Parihar CM, Rana KS, Parihar MD. Crop productivity, quality and nutrient uptake of pearl millet (*Pennisetum glaucum*) and Indian mustard (*Brassica juncea*) cropping system as influenced by land configuration and residual effect of nutrient management. Indian Journal of

- Agricultural Sciences. 2009;79(11):927-930.
14. Sharma P. Response of mustard (*Brassica juncea*) varieties to fertility levels in alluvial soils of Madhya Pradesh. A Journal of Multi-disciplinary Advance Research. 2013;2(2):47-49.
  15. Singh T, Rana KS, Satapathy BS. Productivity, compatibility and economics of mustard (*Brassica juncea*) and (*Lens esculenta*) intercropping system as influenced by moisture conservation practices and fertility levels under rainfed conditions. Indian Journal of Agricultural Sciences. 2014;84(6):770-773.
  16. Snell PD, Snell GT. Colorimetric methods of analysis, 3<sup>rd</sup> Edn. Vol. II D. Nostrand Company, Inc., New York; c1949.
  17. Trivedi SK, Pachauri RK, Sharma G, Joshi BS, Rajput B. Effect of moisture Regimes, NPK and zinc levels on growth, yield, quality, nutrient uptake and economics of mustard (*Brassica juncea*). Journal of Soils and Crops. 2013;23(1):78-85.
  18. Yadav RK, Shivran AC, Jakhar P, Jagetiya A, Puniya MM. Performance of rainfed moth bean and sesame as influenced by intercropping systems and sulphur fertilization. Annals of Arid Zone. 2014;53(2):87-90.